INDEX

Prepared by DR. J. BEN HILL of the Botany Department, THE PENNSYLVANIA STATE COLLEGE, State College, Pennsylvania and Mrs. HELEN DEUSS HILL. Page numbers printed in black-faced type refer to headings of articles or of sections. Numbers in italics indicate references included in lists of "Literature cited."

Acer Pseudoplatanus 569
ADKINSON, JUNE 576, *581*
Alcohol and fertility of white rats 117–141
Alcohol
  As selective agent 134
  Effect of genetically 139–141
  Eliminating germ cells or zygotes 140
  Reduces number of young 139
Alcoholism and the growth of white rats 427–444
  Administration of alcohol by inhalation 428
  Alcohol as selective agent upon genetic differences 428
  Alcohol effect of fumes on treated animals and descendants 427
  Alcohol effect on growth of rats, 438, 440, 441, 444
  Alcohol effect on weight 442-443
  Alcohol prevents reproduction 444
  Alcohol reduces rate of growth 440, 441, 444
  Alcohol treatment retards growth of rats 438
  Differences between weights of tests and controls 433
  Effects of alcohol 443
  Frequency distributions of weights of treated males at different ages 436
  Growth curves of test and control females 437
  Methods of experimentation 428–431
  Offspring of treated rats genetically superior to controls 441
  Offspring of treated rats heavier than controls 439, 440, 444
  Offspring of treated rats more fecund than controls 431
  Rats, growth retarded by alcohol treatment 438
  Selective effect of alcohol on rats 440, 441
  Treated females less often pregnant than controls 431
  Treated males grow less rapidly than controls 435
  Treated rats less fecund than controls 431
  Weight data from treated males 435
  Weight per female rat 432–433
  Weight per male rat 430–432
ALDER, B 274, 275, 277, *317*
ALDROVANDUS 466
ALLEN, E 517, *550*
Allergic conditions 581
Alliogamous population, selection in **508**–512
ALTENBURG, E. 386, *425*, 597, *601*
Andromonoecious types of maize **203**–236
Angioneurotic Edema **573**–582
  Anaphylaxis 579
  Causes 573–574
  Cross of affected×unaffected 577
  Description 573, 576
  Edema of the glottis 577
  Factors possibly two in inheritance 580
  Familial **577**–581
  Heritable 574
  Inheritance explanation 580
  Parents neither affected 579
  Sporadic **575**
ANONYMOUS 466, *506*
Anther ear type of maize **203**–236
Aphis avenae, selection ineffective 143
Arcella 144, 176, 181
Arenga saccharifera 150
ARLITT, A. H. 428, 441, *444*
Ascomycetes 149
Ascus stage of fungi imperfecti 149
Assortment, independent 397
Asthma 575, 581
  Heredity 576
  Recessive 576

BABCOCK, E. B. 338, 535, *550*
BALL, E. D. 274, 275, 277, *317*
BALLY, W. 514, *550*
BAILEY, I. W. 517
BAILLY-MAITRE, J. 467, 473, *506*
BARBER, M. A. *182*
Bar-eye a sex-linked character 5
Bar-eye not fully dominant 5
Bar-eye, origin in *Drosophila melanogaster* 5
BARROWS, WILLIAM MORTON **573**–582
Bars in pigeons **466**–506
BARTLETT, H. H. 146, 147, *183*, 545, *552*
BATESON, W. 545, *550*
BAUR, P. 333
BELLING, J. 526, 528, 532, 545, *550*
BENEDICT, F. G. 272, *273*
Bigenic character, determination of loci 407
Bigenic hypothesis 387, 391, 392, **397**–400
Bigenic nature of ski in *D. melanogaster* **391**–394, 399
Biometric studies of variable groups 182
Black in pigeons **466**–506
BLACKBURN, K. B. 532, *550*
BLAKEMAN, J. 280, *317*
Blakeslee, A. F. 275, 302, 303, 307, 308, *317*, 446, 447, 451, 452, 455, 463–*465*
BLAKINGHEM, L. 203, *226*
BOITARD 467, 498, *506*

Bolten, G. C. 577, *581*
Bonhote, J. L. 468, 475–477, *506*
Bouin (solution) 517
Boveri 523
Brachydactyly 585
Bravais 259
Bregger, T. 597, *601*
Breitenbecher, J. K. 484, *507*
Bridges, C. B. *334*, 338, 344, 355, 373, 376, *379*, *380*, 386, 389, 391, 394, 396, 397, 411, 412, 418, 419, *425*, 527, *550*
British Museum 467
Brown, Aaron 575, *581*
Bruchus 484
Bud mutations associated with tetraploidy 543
Buffon, Count de 466, *507*
Bulloch, W. 577, *581*
Bursa 530
Bussey Institution of Harvard Univeristy 237, 516

California, University of 389
Calliphora erythrocephala 588
Cameron, C. 577, *581*
Canna 532
Carbohydrate, a rare, in waxy maize **568**–572
Card, L. E. 317, 447 452, 464, *465*
Castle, W. E. 543
Cataract 583
  In female line 584
Cattle dominant white 424
  Recessive albino condition 424
Causation and path coefficients **258**
Cell division in wheat species 544, 548
Centropyxis 144, 176, 181
Chapman, E. R. B. 472, 498, *507*
Characters arise by recombination 422
Characters, similar, conditioned by different genes 424
Checks in pigeons **466**–506
Chelidonium majus 569
Chromosome
  Abnormal and sterility 534, 540
  Behavior and sterility 533–540
  Behavior in partially sterile hybrids (see wheat) **513**–552
  Complex in Drosophila species 337
  Duplication 526
  Groups of, in various species of Drosophila 336
  Number and cell size correlated 519, 523, 540
  Number and size of pollen grains in wheat species 523–526, 549
  Number and sterility 534
  Number, doubling results in duplication of genetic factors 526–527
  Number in wheat 514–515, 525–526, 540–543, 547, 557
  Numbers in plants 526
  Ratios influence characters other than sex 527
  Rearrangement in Drosophila species 342
  X in *Drosophila obscura* longer than in other species 374
Chromosomes
  And linkage groups 341
  Assortment in polyploid species 527–530
  Behavior in reduction divisions of species of Rosa 531
  Complete set necessary for gametic and somatic development 537
  Five pairs in *Drosophila obscura* 341
  In cultivated species of wheat **517**–521
  Of rye 519
  Reduplication does not cause sterility 534
  X of three Drosophila species, maps 375
  X parallelism 337
Chrysanthemum chromosome number 527, 545
Clausen, R. E. **385**–426, 424, *426*
Coca, Arthur F. 573, *581*
Cole, L. J. 468, 472, 473, 480, 502, 503, *507*
Collins, G. N. 568, 571, *572*
Collins, J. L. **385**–426, 535, *550*
Color in flowers effect of temperature on inheritance 333
Color patterns in doves corresponding to those in pigeons 502
Columba (see pigeons)
Columba affinis 471, 505
Columba guinea, 467, 502
Columba gymnocyclus 467, 505
Columba intermedia 494, 495, 505
Columba leuconata 467
Columba livia 467, 468, 494–496, 500, 502, 505
Columba oenas 472
Columba rupestris 467, 505
Columba schimperi 467, 505
Columbae 468
Columbia University 389, 391, 394, 396
Columbidae 467, **501**–502
Columbigallina rufipennis 502
Compound, defined 345
Connecticut Experiment Station 237–239, 255
Cooke, Robert A. 574, 576, *581*
Corbié 467, 498, *506*
Corn (see maize)
Cornell University 203, 559, 597
Correlation and path coefficients **258**–273
Correlation in hybrid sterility in wheat 514
Crepis chromosome number 526, 545
Crepis Reutreiana "*gigas*" *mutant* 526
Crepis tectorum×*Crepis virens* 535
Crepis virens 526, 535
Cromwell, Hobart *334*
Crossing over 373
  Correlated with cytological length of chromosome 374
  In Drosophila and change of temperature 338
  In maize **597**–601
  In sex-linked groups in *Drosophila obscura* 340–341
Crowder, Joseph Reed 577, 578, *581*
Crowder, T. R. 577, 578, *581*
Culms inheritance in maize **559**–567
Cunier 583
Cytological methods in wheat 517

DAFERT 569
Daltonism 583, 585, 588
DARWIN, CHARLES 467, 468, 471, 473, 480, 487, 495, 505, *507*
Datura, chromosome number 526, 528, 530, 532, 533
  New species 534
  Normal 534
  Poinsettia mutant 528
  Sterility 533
  Triploid 534
  Trisomic 534
DAVENPORT, C. B. 585, *589*
DAVIS, B. M. 530, 531, *550*, **590**–*596*, 591, 595
Deficiency in *Drosophila obscura* 355
DETLEFSEN, J. A. 424, *426*
Dextrins 568
Diandric heredity WZ type of sex linkage 583
Diagynic heredity XV type of sex linkage 583, 589
Didynium nasutum 144, 176, 181
Difflugia 144, 149, 157, 178, 181
DIGBY, L. 526, *550*
Diptera 372
Disease resistant races in maize 252
Dominance reversed 527–528
DORSEY, M. J. 542, *550*
Doves and pigeons, similar color patterns 502
Doves, species of 502
Drosera 531, 532
Drosophila 349
  Graded multiple-allelomorph series 504
  Intersex 527
  Interspecific crosses 376
Drosophila ampelophila 337
Drosophila funebris 351, 376
Drosophila hydei **319**–334
  Backcross ratios 324
  Backcross 325
  Combination of scarlet and variable under different temperatures **326**–330
  Crosses
    Backcross hybrid ♀ × variable scarlet ♂ 331
    Backcross variable scarlet ♀ × hybrid ♂ 331
    F_2 from normal-eyed variable scarlet ♀ × wild stock ♂ 329
    F_2 from wild stock ♀ × normal ♂ from variable 330
    F_2 from wild stock ♀ × speck-eyed scarlet ♂ 329
    F_2 from wild stock ♀ × variable scarlet ♂ with minimal eyes 329
    Normal-eyed ♀ from variable scarlet stock × wild stock ♂ 328
    Normal-eyed scarlet ♀ from pure minimal mass culture × wild stock ♂ 330
    Normal mahogany ♀ × variable scarlet ♂ 331
    Variable scarlet ♀ × normal mahogany ♂ 331
    Wild stock ♀ × minimal-eyed scarlet ♂ 329
    Wild stock ♀ × normal ♂ from minimal 330
    Wild stock ♀ × scarlet variable ♂ with normal eye 328
    Wild stock ♀ × variable scarlet with very small eyes 328
  Double recessive scarlet variable 324
  Environmental influence on mutant 325
  Environmental stimulus causing phenotypic differences 333
  Eye mosaics **333**
    Inheritance effect of environment 333
  Linkage 320
    Of variable and scarlet 319–320, **323**–325 331, 334
  Mahogany normal 331, 332
  Mahogany variable 331, 332
  Mosaic 333
  Mosaic male 333
  Mutant dependent on environment 319 322–330
  Mutant scarlet 321, 333
  Mutant "variable" 319
  No crossing over in the male 331
  Scarlet 320, 324, 330
    A new mutant eye color 320
    Normal 331, 332
    Variable 322, 325, 331, 332
      In reduced temperature 328
      In warm temperature 327
  Sex ratio 325
  Variable 324, 325, 333
    Behavior in inheritance **322**–323
    Dependent for expression on environment 334
    Is recessive 320
    Origin and description **321**–322
    Response to temperature 332, 333
    Reversion due to low temperature 332–333
    Scarlet 325
    Stock cultures 330–333
    Stock established by selecting variable eye **330**, 332
    Under controlled temperature 319, **323**
    Wing mutation in culture variable 330
Drosophila melanogaster 333, 337–340, 342, 345, 347, 351, 354, 355, 357, 359–361, 367–369, 374–379
  Abnormal abdomen 333
  Bar 391
  Bar-eyed mutant expression of temperature 333
  Black 397, 407
  California curled 389
  Club 388
  Curled 388, 389, 390
  Dichaete 386, 391, 397–399, 401, 402, 413–415
  Dominant mutant genes 400
  Duplex 413, 414
  Duplication 386
  Eosin eye color modifiers 386
  Eye color modifiers 504

Drosophila melanogaster (continued)
Eye facet number expression of temperature 333
Genes, modifying 386
Inheritance of ski wings 385–426
Jaunty 388–390
Lethals balanced 386
Mutation by recombination 423
Mutation, point 423
Non-disjunction 386
Notch wings 386
Ratios wild type to ski 403–407
Sable 527
Scute 376
Selection in White Bar Eye 1–115
Bar eye 2, 3
Changing to full eye and ultra-bar 4
Coefficient of dominance 21
Facet number affected by temperature 3, 8
Factorial units 5
Low bar 22, 23
Not fully dominant 5
Origin 5
Red bar stock 7
Selection for high facet number 3, 114
Selections for low facet number 3, 114
Sex-linked character 5
Sex ratios affected by lethal mutant factors 70
Sexual dimorphism 6–7, 92
Change in 70
Stock 2, 5–7
Germinal diversity 2
Ultra-bar 16, 22, 114
Vestigial-winged bar 3, 7
White bar-eye race 1–115, 7
Class facet values 15, 20
Correlation between degree of selection and effectiveness 107–112
Crosses
Between high and low lines 3
Ultra-bar × bar 24, 28
Dominance of bar and full 21
Facet and factorial values of selected pairs for all matings 32–44
Facet differences not of hereditary value 112
Facet factors eliminated by selection 4, 114
Facet number affected by sex-linked lethal factor 86, 113–115
Facet number conditioned by elimination of factors 86
Facet number factors environmental effect 14–16
Facet number factors germinal effect 14–16
Facet number inhibitor 16
Facet number 26
Factorial values for facet number 13, 15, 20
Factorial values frequencies 54–69, 71–78
Formula for coefficient of dominance of bar-eye 21
Full eye 4, 22–23, 112, 114
Facet number 5
Germinal differences 84, 88
Arise by mutations independently of selection 113, 115
Heterozygosity effect in bar eye 84
Low-selected bar 16
Means, female and male 92–103
Mutants to full eye and to ultra-bar 49, 112, 114–115
Mutation to full eye from selected bar 84, 112
Mutations independent of selection 114
Non-heritable fluctuations 84
Ontogenetic development of characters 2
Pedigrees in high line 10
Pedigrees in low line 9
Restriction in range, selection in bar eye 84
Rudimentary 5
Selected lines of bar-eye range 78–87
Selection
Bar eye matings in 8
Bar eye original data 24
Effect due to elimination of germinal groups 78–87, 113
Effect on females 45, 113
Effectiveness 112, 113
Difference between parent and offspring 95–98
Eliminating facet factors 4
For eye facet number 1–115
Germinal changes 2
In bar eye temperature control 8
Lines, low and high compared 24, 25
Methods 7–12
On basis of performance 88
Positive results by conserving mutation 4
White bar 12, 19–21, 26
High selection 27, 30, 47–48, 52–69, 91, 92, 114
Low and high lines contributions to divergence 91–92
Low and high lines differences 88–91
Low selection 26, 27, 29, 45–46, 50–51, 54–69, 92, 114
Selections eliminating germinal factors 112
Sex-linked lethal factor 92
Standard deviation 99, 104–106
Synchrony between low and high generations 12
Temperature effect 16
Ultra-bar mutants 85, 112–114, 115
Unselected bar 16
Variability in ultra-bar bar and full eye 22
Sepia 393, 397, 405
Sepia locus 396
Sex-linked jaunty 389
Ski 398–400, 402, 407
Ski-II 385
Determination of locus 407–412
Ski-III 385

Drosophila melanogaster (continued)
Determination of locus **413–420**
Ski **385–426**
Bigenic character 387, 399, 407, 425
Bigenic hypothesis 387, 391, 392, 393, **397–400**
Bigenic nature 388, **391**–394, 399
Characters employed in analysis **396**–397
Chromosome position 393
"Crimp" in costa or marginal vein 390
Description **389–390**
Differs from wild type in two pairs of genes 400
Duplex 389, 392, 393, 398, **400–403**, 413
Examination of flies 396
Experimental methods **394**–396
For instructional purposes 421
Genotypes 392
Inheritance 424
Linkage relations 393
Locus 388
Maintaining stocks 395
Mode of inheritance 424
Not a monogenic character 399
Origin and history **387–389**
Procedure with experimental cultures 395
Ratios wild to ski **403–407**
Recombination in connection with origin of 422
Resembles jaunty and curled 388, 389
Reversion to wild type 388
Segregation 393
Simplex 392, 394, 400–403, 404
Morphological differences from duplex flies 403
Mated *inter se* 403, 404
Phenotypically identical with jaunty 389
Star ski 398–401, 407, 414
Technique of breeding 394–396
Types of reciprocal matings 392
Variable in expression 389
Spineless 397, 413
Spines number 386
Star 391, 396–402, 407, 414
Star black 407
Star dichaete 397–402, 413
Streak 386
Truncate wings 386
Vermillion 377, 527
Vestigial modified by differences in temperature 333
Vortex 386
White 358
White-eyed males 387
Wild type constant 385–386
Drosophila mulleri 321
Drosophila obscura **335–384**, 387
Allelomorphic series, white and eosin multiple 359
Autosomal characters 357
Bithorax 357
Chromosome
Genetic length 340
Localization of genes **348**
Long 341
Map **373–374**
Involving beaded, vermilion, fused and short 349
Rearrangement 342
X, loci in 384
X very long 379
Chromosomes, five pairs 341
Crosses
Glazed×beaded **364**
Glazed×pointed **364**
Miniature×fused 368
Miniature×vermilion 368
Notch×reduced rough 354
Pointed×beaded 362, 363
Polished×rough 371
Reduced rough×vermilion 350, 371
Singed male×fused short female 369
Singed×yellow vermilion 370
Vermilion×beaded 348
Vermilion×short 348
Crossovers between
Ascute with pointed and yellow 367
Beaded-vermilion-fused and short 348
Compressed and fused 367
Glazed and beaded 364
Miniature and reduced 369
Notch and reduced 354
Notch and shorter 352
Reduced and rough 351
Scutellar and yellow 361
Vermilion and broken 365
Vermilion and reduced 354, 369
Vermilion beaded and short 348
White and fused 370
White and notch 358
White and singed 369
Crossing over between sex-linked characters **339–341**
Crossing over in X chromosome 379
Culture methods **338**
Deficiency 355
Difference in sequence 360
Dominance relations of short allelomorphs 344
Genes, order 348
Genes, sequence 351, 359, 360, 379
Lethal factors 372
Lethal, notch 351
Linkage experiments 382–383
Linkage groups 339, 341
Linkage of
Compressed and fused 367
Dwarf and short 371
Glazed and fused 363
Rough and fused 351
White and yellow to notch 358
Yellow and notch to eosin 359
Linkage relations of sex-linked characters **335–384**
Loci of reduced and rough **350–351**
Loci of yellow, notch and white or eosin **359–360**

Drosophila melanagaster (continued)
Modifying factor 335, **342–345**
Multiple allelomorphic locus 335, **342–345**
Mutability 335, **339**
Mutants
Achaete modifier 335, **355**
Achaete notch 356
Ascute 336, 367, 378, 381
Between fused and short 367
Locus 367
Beaded 335, **345**, 347, 361, 362, 363, 373, 381
Locus and valuation **346**, 374
Broken 336, 347, **364**, 381
Dominant in females 364, 365
Lethal in male 364
Locus **365**
Sex-linked recessive lethal 365
Compressed 336, 350, **366**, 374, **378**, 381
Locus **367**
Deformed 336, **371**, 381
Lacking in vigor 372
Dwarf 336, **371**, 381
Eosin 336, 342, 354, **358**, **359**, 376, 381
Extended 336, **372**, 381
Lacking in vigor 372
Fused 335, **346**, 347, 363, 370, 373, 381
Locus 374
Gap 335, **344**
A modifier of short 344–345
Glazed 336, 350, **363**, **377**, 381
Homozygous females sterile 363
Locus **363**
Grooveless 367
Lethals 381
Miniature 336, **368**, 370, 371, **377**, 381
Locus **368**
Minus, autosomal 349
Minute 356
Notch 335, 342, **351**, **352**, 361, 381
Deficiency nature **354**
Locus **353–354**
Recessive lethal 351
Similarity to notch mutants in other species 351
Specific notch modifiers **355**
Notch$_2$ 335, **352**, 355
Notch$_3$ 335, **353**
Pointed 336, **361**, 362, 367, 381
Locus **362**
Polished 336, **370**, 371, 381
Locus **370**
Reduced 335, **349**, 350, 354, 371 381
Rough 335, **350**, 354, 381
Nature and cause of character 350
Scutellar 336, 355, **360**, 376, 381
Locus **361**
Short 335, **342**, 347, 363, 381
Shorter 335, **343**, 381
Shortest 335, **343**, 381
Singed 336, **369**, 371, **377**, 381
Females sterile 369
Locus 369
Slender 336, **365**, 381
Locus **366**
Split 356
Modifier of notch 335, **356**
Notch$_2$ 356
Stubby 336, **372**, 381
Vermilion 335, **347**, 362, 364, 370, 371, 374, **377**, 381
White 335, 342, 347, 353, 354, **357**, **359**, 369, 370, **376**, 381
White$_2$ 336, **357**
Yellow 335, 342, 353–357, 361–363, 367, 370, 371, 373, 374, **376**, 381
Notch combination 353
Mutants similar to those of other species and corresponding chromosome regions **375**
Mutation, frequent in middle region of X chromosome 347
Mutations identical with those in other species 339, 379
Mutations in X chromosome 378
Non-disjunction of sex chromosomes not found 339, 340
Scarlet 357
Sex-linked characters description and location **342–373**
Sex-linked group compared with other species of Drosophila 373
Slender-bristle stock 356
Vermilion and broken crossovers 365
Vermilion and reduced Broken crossover 365
X chromosome includes a region not present in X chromosome of *D. melanogaster* 339
Drosophila repleta **319**, 320, 321
Eye color mahogany 322
Seven modifiers acting on eye color 504
Sex determined by ratio of X chromosomes to autosomes 527
Sex-linked mutant 320
Drosophila simulans 337, 347, 377
Sterile F_1 535
Drosophila species forked 372
Drosophila superiority for genetic investigation 425
Drosophila virilis 337, 347, 374, 375, 377–379
Drosophila willistoni 377
Drosophilae 321
DUDLEY, A. H. 514, *550*
Dwarf type of maize **203–236**

EAST, E. M. 203, 219, *226*, 237–240, 248, 250–252, 254, *257*, 514, 517, 527, 539, 540, 542, 545, *550*, *551*
EATON, J. M. 467, *507*
Eczema 575, 581
Edema of the glottis 573, 577
EDGERLY, EDWARD T. 577, 579, *581*
EGBERT, A. D. 274, 275, 277, *317*
Egg Production in fowl **274–318**
Egg production in Rhode Island Red breed of domestic fowl **446–465**
Biological factors on which fecundity depends **447**

Egg production in Rhode Island Red breed of dometic fowl (*continued*)
Correlation between annual production of deviation of individual months **455–457**, 464
Correlation between individual months and annual production **448–454**, **455–457**, 464
Correlation between records of individual months of same year **457–463**, 464
Correlation of individual month and other eleven months **454**, 455
Correlations of egg production of Rhode Island Red agree with those for White Leghorns 465
Distribution of magnitudes of correlations between individual months and
Antecedent months 461
Subsequent months 460
Physiological activities at various periods interrelated 446
Regression equations for prediction of annual egg production 450
Regression of annual production on monthly production 451–453
ELDERTON, W. P. *317*
EMERSON, R. A. **203–236**, 210, 219, *226*, 559, 560, *567*, 597, 598, *601*
EMERSON, STERLING H. **203–236**
Endosperm development and F_2 sterility in wheat **553–558**
Endosperm formation not due to triploid condition 558
ENRIQUES, PAOLA **583–589**
Epistatic series in pigeons **480–487**, 499, 500, 502, 503
Erythrodextrin 568–570
Occurrence in plants 569–570
Evolution
By hybridization 422–424
By mutation 182
By selections of fluctuating variations 182
EWING, H. E. 143, *182*
Eye color, mahogany in wild type *Drosophila repleta* 322
Eye selection 330, 332
Eyeless mutant in *Drosophila hydei* **319–334**
EYSTER, W. H. **559–*567*, 597–*601***

Factors presence and absence theory 424
FARNHAM, M. E. 526, 528, *550*
Fecundity dependence on biological factors 447
FEDERLEY, H. 532, *550*
Fertility of rats treated with alcohol 440
Fertility of white rats, influence of alcohol **117–141**
Fertility reduced by alcohol treatment 139
FIELD COLUMBIAN MUSEUM 467, 506
Fleming solution 517
Fowl, correlation between first and second year egg production **274–318**
Annual egg production 307
Anatomical fecundity 276
Autumn production 299, 302
Barred Plymouth Rock 276
Color criteria in egg production 275
Condition of birds with respect to laying activity 302
Correlations between
Annual production for two years 281
Egg production of first and second year **274–318**, 276, 280, 283, 294, 305, 313, 314, 316
First and third year 277
Individual months of first two years 290, 291
Individual months of first and second year 285, 287, 294, 296
Individual months of second year and first year 286
Inter-annual 311
Inter-annual for monthly production **287–293**, 296, 297
Genetic significance **293–302**
Inter-annual for "cycles" of first and second year 299
Inter-annual for various months of first and second year 284
Inter-annual of flock, means **283–284**
Monthly records of one year and annual record of another year **284–287**
Second and third year 277
Summer months of first year and winter months of second year, low 298
Summer months of first year and months of various periods of second year 298
Winter months of two years 294
Correlation, intensity between two years 301
Correlation significance of deviation 301
Correlations, summary of data 300
Criteria in egg production 275
Cycles 296, 297
Egg production
Color criteria 275
First and second year 308–310
Inter-annual correlation 275, **278–280**
Primary factor 282
Mean annual 281, 306
Seasonal distribution 277
Fecundity of diverse periods 275
Fertility dependent on Mendelian factors 293
First year production, mean 278, 303, 304
Individuality of the bird 282
Inheritance of fecundity 274, 275
Linearity of regression in egg production 280, 286
Mean production of first and second years correlated 314, 315
Mendelian factors for winter production 293
Not differentiated by summer and autumn production 300
Production lower second year than first year 305
Ratios of correlation coefficients to their probable errors 290, 292
Records by months 311–313
Records, egg production of first and final

Fowls (*continued*)
months of first year related to second year's record 310
Regression equation in egg production 279, 287
Regression linearity in egg production 280, 286
Regression of second-year annual production on first year monthly production 288, 289
Second-year egg production, mean 278, 279, 303–305
Selection, basis 305
Spring production 299
Spring or reproductive cycle 293, 294, 300
STORRS INTERNATIONAL EGG-LAYING CONTEST 302–308
Summer production 299
VINELAND INTERNATIONAL EGG-LAYING AND BREEDING CONTEST 277, 282, 304–308, 314
White Leghorns 276, 277, 302, 316
Heritable differences 300
Winter cycle of egg production 293, 294, 300
Winter production of first and second year 293
Winter production 299, 302
Fowl
Domestic, egg production 446–465
Treated with alcohol, growth curves 443
Fowls
Domestic, Silky, White Leghorns, White Plymouth Rock, White Wyandotte, 424
Treated with alcohol and ether, fertility data 137, 138
Whites dominant 424
Whites recessive 424
FREUDENTHAL, WOLFF, 576, *582*
Fungi Imperfecti 142, 145, 149

GAINES, E. F. 530, 535, 538, *551*
Gallus (see fowl)
GALTON, FRANCIS 259, *273*
Gametophytic characters controlled by sporophyte 547
Gametophytic chromosomes control of characters by 547
GARBER, R. J. 238, 239, 251, *257*
GATES, R. R. 527, 530, 531, 534, *551*
GAUSS 259
GEERTS, J. M. 531, *551*
Gene, change in composition 424
Genes, different conditioning similar characters 424
Genetic differences, effect of alcohol on 428
Genetics of maize 237, 238
Gentiana lutea 569
Geopelia striata 502
Germ-cells eliminated by alcohol 134, 140, 141
Germ-cells inferior eliminated by alcohol 134
Germinal differences in *Drosophila melanogaster* 84
Germinal modification due to alcohol 139, 140
GHIGI, A. 505, *507*
GIBSON, W. 280, *317*
GOODALE, H. D. 446–*465*, 447
GOODSPEED, T. H. 424, *426*
GOWELL, G. M. 275, 276, *317*
GOWEN, J. W. 419, *426*, 597, *601*
GRANDIDIER 583, 585
GREGORY, R. P. 528, *551*
GRIFFEE, FRED 239
GRIS 569
Growth factors in maize 220, 221
Guinea pig 264–268
Birth weight 265, 266
Drunken by inhalation 123
Fatally affected by large doses of alcohol 136
Genetically selective action of alcohol 139
Gestation period 264, 266
Litter size 126, 264–266
Prenatal growth curve 265
Rate of growth 266
Reduction of litters by alcohol 135

HAGEDOORN, A. L. 422, *426*
HAGEDOORN, VORSTHEUVEL LA BRAND, A. C. 422, 426
Haidenhain (iron-haemotoxylin) 517
HANEL, ELISE 143, *182*
HANNAS, RALSTON R. 277, *317*
HARPER, R. A. 571, *572*
HARRIS, J. ARTHUR 143, *183*, 270, 272, *273*, 274–318, 275, 281, 301–303, *317*, 446–465, 451, 452, 455, 457, 463, 464, *465*
HARRISON, J. W. H. 532, *550*
Hay-fever 575, 581
HAYES, H. K. 203, *226*, 237–*257*, 238, 251, 527 536, *551*
HEGNER, R. W. 144, 176–179, 181–*183*
Helminthosporium 182
Hemerolopia 588
Hemophilia 583, 585, 588
In the female line 584
Heredity
Hologynic, a new type 583, 589
Of angioneurotic edema 573–582
Heritable characters, methods of modifying and controlling 427
Heterozygosity effect on bar eye in Drosophila 84
Hevea brasiliensis 168
Hexaploid forms 526
Hexaploidy in wheat species, origin of 540–541
Hieracium auricula×*H. aurantiacum* 531
Hieracium chromosome number 526, 532
Hieracium excellens 531
High protein maize produced by Mendelian methods 237–257
VAN'T HOFF 16
HOGE, MILDRED A. 333, *334*
Holoandric heredity, through male line 583, 589
Hologynic heredity 583–589
Explanation 586–587
Hemophilia 584
Heredity through female line 583
Known only in man 589
Selective fertilization 587, 588
Senile cataract 584
Sex-linked characters 583

HOWARD, A. 530, *551*
HOWARD, G. L. C. 530, *551*
HORTON 254, 255
Human basal metabolism 272
  Body surface 272
  Body weight 272
  Correlation coefficients 272
  Heat produced 272
  Hereditary and environmental factors 272
  Path coefficients 272
  Stature determines body surface 272
  Stature determines body weight 272
HUTCHISON, C. B. 598, *601*
Hybrid vigor effect 533
Hybrids, sterility in wheat **553**–558
HYDE, ROSCOE R. **319**–334, 320, *334*
Hydra selection not effective 143

INDIANA UNIVERSITY 568
Inheritance
  Modified by temperature in Drosophila 333
  Of fecundity in fowl 274
  Of ski wings in *Drosophila melanogaster* **385**–426
  Of zigzag culms in maize **559**–567
INTERNATIONAL EGG LAYING CONTEST 448, 451
Iris germanica 569

JENKINS 239
JENNINGS, H. S. 142, 144, 157, 174, 178, 179, 181–*183*
JOHANNSEN, W. 142, 143, 175, 181, *183*
JOHNS HOPKINS UNIVERSITY 258
JONES, D. F. 238–240, 248, 250–252, 254, 255, *257*, 571, *572*
JONES, SARAH VAN HOOSEN **466**–506

KAUFFMAN, C. H. 150, *183*
KELLEY, FRANK J. 469, 486, *507*
KEMPTON, J. H. 203, *226*, 569, 570, 571, *572*
KIHARA, H. 514, 515, 537, 539, 540, *551*
KIRKPATRICK, W. F. 275, 302, 303, 307, 308 *317*, 446, 447, 451, 452, 455, 463, *465*
KÖRNICKE, M. 514, *551*
KRAFKA, JOSEPH, JR. 1, 3, 8, 12, 13, 16, 17, *115*, 333, *334*
KURTZWEIL, C. 536, *551*

LARUE, CARL D. **142**–201, 146, 147, 168, *183*
LANCEFIELD, D. E. **335**–384, 345, 372, *379*
LASHLEY, K. S. 143, 144, *183*
Lepidoptera 587
Lethal factors in *D. obscura* 372
LEWIS, HARRY R. **274**–318, 277, 303, *317*, 447, 457, *465*
LIEBIG 168
LINDSTROM E. W. 560, 565, *567*
Linkage in *Drosophila hydei* 320
Linkage in maize 560, **565**, **597**–601
Linkage relations in *Drosophila obscura* **335**–384
Litters, number compared in guinea-pig and rat treated with alcohol 137
LOTSY, J. P. 422, *426*

LUSK, FRANK B. 577, 579, *581*
LUTZ, A. M. 526, *551*
LYELL, J. C. 473, 487, *507*

MACDOWELL, E. C. **117**–*141*, 122, **427**–445, 428, 431, 435, 440, *444*
MCCLUNG 517
MAINE AGRICULTURAL EXPERIMENT STATION 516
Maize
  Aleurone color 424
  Andromonoecious type recessive to monoecious 203
  Andromonoecious types **203**–236, 225
  Anther ear type **203**–236, 207, 222, **224**–225
    Characteristics **204**–210
    Double recessive 207–210
    Inheritance of **212**
    Larger 208
    Seeds 206
    Simple recessive to normal A_nA_n 212, 225
    Small 208
  Breeding 238
    For economic characters 257
    Literature 238
    Materials and methods **239**
  Carbohydrate in waxy maize **568**–572
  Chlorophyll degeneracies 252
  Corn smut, 252
  Crop improvement problems 237
  Cross-pollination effect 239
  Crosses 212–236, 561, 566
  Crossing over **597**–601, 598
  Crossing over, frequency in microspore and megaspore **597**–601
  Dextrins chemical tests for 568, 569
  Dextrins rare in tissues 568
  Dominance of low protein over high 251
  Dwarf anther ear 225
    Description 207–210
    Double recessive 224
    Infertility 210, 225
    Mature plant 210
    Simple recessive to normal DD 211, 225
  Dwarf type **203**–236, 207, 208, 224, 225
    Characteristics **204**–210
    Height 218
    Inheritance **210**–212
    Seeds 206
  Ear types 205
  Endosperm 527, 570
  Endosperm of waxy maize, erythrodextrin 568–570
  Factors 218, 220, 225, 559, 560, 567
  Genetic interrelation of dwarf and anther ear types **212**–218
  Genetics 237, 571
    Protein content 238
  Growth factors 220–222, 226
  Height segregation 224
  Height 223
  Heterotic effect of cross-pollination on protein content 248, 251
  High and low races 237

Maize (*continued*)
High protein races 237, 238, 242, 253
High protein×low protein F_1 249
Inheritance of protein content **251–252**
Inheritance of zigzag culms **559–567**, **560–565**
Linkage 225, 226, **565–567**, 597–601
Low protein races 237, 248
Mass breeding, difficulties in technique 239
Mode of growth 560
Multiple factors affecting height growth 221
Normal 207, 224, 225
Heights 218
Oil content high and low races 237
Perfect flowers 203, 225
Pre-Columbian occurrence in Asia 568, 572
Production of high protein **237–257**
Protein content 239, 240, 241, 252–254, 256
Analyses 239
And cultural conditions **239–242**
And number of seeds per ear **242–248**
And yields compared 257
Comparison in self-pollinated and wind-pollinated **248–251**
Crosses between high and medium protein races 251
High and low races 237, 238, 242, 253
Of self-fertilized ears and protein in progeny 241, 242
Yields and shelling percent compared 256
Pseudo-starchy endosperm 571
Ratio modified 561, 567
Root rot 252
Rudimentary stamens **224–225**
Seedlings of various types 209
Selection experiments 237
Selection possibilities 237
Self-fertilization and isolating races for protein content 251, **252**, 253
Self-fertilized lines and use in improvement **252–257**
Self-fertilized lines decrease in yield 250
Self-fertilized lines increase of protein content 251, 254
Self-fertilized strains and resistance to plant diseases 252
Semi-dwarf andromonoecious 203
Strains
Illinois high protein 255
King Phillip Flint 240, 242, 247, 249, 253
Leaming 247, 255
Longfellow 249
Mercer Flint 243, 247, 249, 253
Minnesota No. 13 240, 241, 243, 244, 247–251, 253–257
Missouri dent corn 559, 566
Red Flint 240, 247, 250, 251
Rustler dent 240, 244, 247, 249, 253
Studies reviewed **238**
Variety, a synthetically produced 255, 256
Tall normal 208, 219
Tom Thumb pop corn 208, 218–220, 222, 559, 566
Tom Thumb ear type 205
Tom Thumb mature plants 208
Variability in stature **218–224**
Waxy endosperm
Description 568
Inheritance 569
Simple recessive to corneous 570
Waxy maize a mutation 571
White forms, three distinct recessive 424
Zigzag culms **560**, 566
Mass breeding technique 239
MASSACHUSETTS AGRICULTURAL EXPERIMENT STATION 448, 463
MAST, S. O. 144, 176, 181–*183*
Matroclinous inheritance in maize endosperm 527
MATTOON, E. W. 2, 6, 19, *115*
MAY, H. G. 3, 7, 8, *115*
Mendelian inheritance of patterns of pigeons 468
Mendelian methods and production of high-protein maize **237–257**
METZ, C. W. 336–338, 340, 342–344, 372, 377–*379*
MEYER, ARTHUR 569, *572*
MIDDLETON, A. R. 144, 175, 176, *183*
MILL, JOHN STUART 260
MINNESOTA EXPERIMENT STATION 237–239, 251, 257
MISSOURI AGRICULTURAL EXPERIMENT STATION 559
Modifier in *Drosophila obscura* 344
MOHR, O. L. 354, *379*, 386, *425*
Monotropa Hypopytis 569
MONTGOMERY, E. G. 203, *226*
MOORE 466, *507*
MORGAN, T. H. 5, 144, *183*, 333, *334*, 355, 361, 373, 378, *380*, 386, 388, 389, 391, 396, 397, 411, 412, *425*, *426*, 527, *551*
MULLER, H. J. *334*, 373, *380*, 386, *425*, *426*, 528, 530, 545, *551*
Multiple allelomorphic series in Bruchus 484
Multiple factors 530
Mutability of *Drosophila obscura* 335, 339
Mutations
Black and black blue-tail in pigeons 486
By recombination 423
Dependent on environment for development 319, 322–330
Diversity of order of genes in different species 379
Identical in various Drosophila species 339, 341–342, 379
In Drosophila 391
In *Drosophila hydei* **319–334**
In Drosophila species compared 375
In Helminthosporium 182
In lower animals 182
In Pestalozzia 167, 171, 182
In polyploid species 528
In yeasts 182
Method for determining identicity in different Drosophila species 376
Non-disjunctional 144
Point 423

Mutations (*continued*)
Frequency of Drosophila 112
Rate of in Drosophila species 339
Similar in different species of Drosophila 376–378
Somatic 144
Myopia 588

NÄGELI 569
NAKAO, M. 514, *551*
NEUMEISTER 473
Neuritis optica 588
NICE, L. B. 428, 442–*444*
Nicotiana hybrids 540
Sterility and incompatible chromosome combination 540
Nicotiana pollen 545
Nicotiana tobaccum, two genetically distinct red-flowering forms 424
NILES, HENRY E. **258**–273
NILSSON-EHLE, H. 530, *551*
NIXON, CLARA 274, 276, *318*
Non-disjunction 527
Absence in *Drosophila obscura* 339, **340**
Non-heritable fluctuation in *Drosophila melanogaster* 84
NONIDEZ, J. F. 403, *426*
NUTTAL, J. S. W. 476, 507
Nystagmus 588

Oena capensis 502
Oenothera brevistylis **590**–596
Factors for style length and stigma structure of slight variability 596
Fertility not due to heritable factors 590
Seeds, treatment to secure complete germination 592
Selection based on performance 591–595
Selection for style length and fertility **590**–596
Results 594
Sterility 590
Style length not due to heritable factors 590
Oenothera chromosome number 526, 528, 533, 545
Oenothera gigas 526, 528, 534, 545
Nanalla mutants of 528
×*Oelata rubricalyx* 534
Oenothera Lamarckiana 526, 534, 590
×*O. gigas* 531
Oenothera lata×*O. gigas* 531
Oenothera lata-like 534
Oenothera lata rubricalyx 534
×*rubricalyx* segregate 534
Oenothera mutants causes 530
Oenothera rubricalyx×*O. gigas* 534
Oenothera semilata 534
Oenothera sterility 533
In pentaploid species 534
Oestrous cycle lengthened by alcohol treatment 137
OHIO AGRICULTURAL EXPERIMENT STATION 508
OHIO STATE UNIVERSITY 573
OLSON, P. J. 239

Ontogenetic development of characters 2
Orthogenesis in pigeons 468, 484–486, 506
Oryza sativa glutinosa 569
OSGOOD, DR. 506
OSLER, WILLIAM 574, *582*
OVERTON, E. 514, *551*

Panicum miliaceum glutinosum 569
PAPANICOLAOU, G. N. 126, 131, 134, 136, *141*, 428, 442, *445*
Paramecium 142, 144, 149
Selection 143
PARKER, J. H. 536, *551*
PARNELL, F. R. 545, *551*, 569, *572*
PASTEUR INSTITUTE 573
Path coefficients **258**–273
Causation **258**–273 259
Coefficient of correlation 262, 269–271
Coefficient of determination 262, 270, 271
Gaussian curve of error 262
Mathematical measure of causation 258
Mathematics of **263**
Method **261**–263, 273
Test of **268**–273
Normal curve of error 262
Partial correlation 273
PEARL, R. 122, 123, 134, 137–*141*, 274, 276, 297, *318*, 428, 440, 443–*445*
PEARSON, KARL 143, 144, 156, *183*, 259, 260, *273*, 280, 301, *318*, 557, *558*
Pentaploid hybrids sterility in Oenothera, Datura and Triticum 534
PERCIVAL, J. 514, 515, 541–543, *551*
Pestalozzia Guepini **142**–201, 145, 146, 149, 157, 159, 162, 175, 181, 182
Aberrant form 171
Advantages for selection studies **149**
Culture methods 150, 151
Description 145
Heritable characters 151
Intermediate line 154, 161, 184–201
Mutation 171
In length of spore appendages 167
Mutations give rise to lines significantly different 182
Results of selection within pure lines **142**–201
Selecting device for isolating spores 168
Selection according to progeny performance **152**–167, 182
Selection biometric study 166, 188, 201
Selection effect 161, 162, 170
Selection for length of spore appendages **162**–167
Selection for length of spores **158**
Selection method 180
Selection, minus 154, 156, 160, 161, 170, 172–174, 182, 184–201
Selection of spores according to visible characters **167**–174
Selection, plus 154, 156, 160, 161, 170, 172–174, 182, 184–201
Selection totally ineffective 182
Spore appendages length 148

Xestalozzia Guepini (continued)
Spore appendages mean lengths 163–166, 189
Standard deviations 189
Spore length 147, 171, 174
Means 160, 161, 184
Standard deviations 184
Spores variability 151
Variability in number of spore appendages 152
PHILLIPS, JAMES MCILVAINE 573–582, 575, 582
PIETERS, A. J. 150, *183*
Pigeons, inheritance 466–506
Bar dominant to barless 484
Barred 467, 472, 483, 487, 495, 498, 506
Bars produce only bars 477
Black and black blue-tail dominant mutations 486
Black baldheads 500
Black blue-tail 469–471, 480, 482, 483, 485
Black check 485
Black, check and bar genetic relationships 473–480
Black, check and bar phenotypic differences 473
Black dominant to check 475
Black 467, 485
Full black 469, 470, 480, 482
Dominant to barred type 474, 475
Black-tailed birds rump color condition 496
Blacks throw blacks, checks, and bars 477
Blue 466, 467, 469
Blue barless 469, 472, 473, 485
Eichbuhlertaube 473
Feld-Tauben 473
Goldgimpel 473
Homer 473, 482, 484, 506
Ice Pigeon 473
Inheritance 484
Pfaffen 473
Strasser 473
Swallow 473, 484
Swiss Pigeon 473
Blue black-barred 466, 467, 469, 472, 473, 480, 485
Blue corresponds to "gull gray" 472
Blue with black check-marks 467
Bluing series 469–473
Check and bar independent Mendelian inheritance 468
Check and bar relationships 468, 469
Check and bar two types in domestic pigeons 501
Check and black blue-tail distinguished 482
Check breeding behavior 483
Check dominant to bar 476
Check 466, 467, 469, 471, 472, 480, 487, 495, 498–501, 506
Checked and barred reversions 468
Checked and barred due to factor recombinations 468
Checks throw only checks and bars 477
Color modification of depth and sheen in black birds 498
Color patterns in pigeons and doves correspond 502
Color types in pigeons epistatic inheritance 469
Colors due to interaction of factors 501
Columba livia progenitor of domesticated pigeon 468
Crosses 478–481, 484, 487, 492, 493, 500, 501
Dilute series 486
Dun checks 486
Dun silver-tails 486
Dun wing bars 486
Full-duns 486
Silver barless 486
Silvers 486
Sootied silvers 486
Dull-black 497–501
Dun checked 498
Epistatic series 480–486, 487, 499, 500, 502, 503
Factors 474, 480, 482, 486, 494, 497, 506
Fantails 482
Genetic relations of blue types to black 482
Helmet 466
Interaction of factors 469
Jacobin 466
Kitiness 469
Light and uniform colored rump 494–497
Orthogenesis 468, 484–486, 506
Patterns two color relationships 466
Pigeon of Barbary 466
Polyphyletic origin of domesticated pigeon 505
Red and yellow 487
Red-barred 487
Red-checked 487
Restricted reds 487
Reversionary blues 471, 480
Reversions 468
Rock-Fantail 472
Rock Pigeon 471, 486, 487, 505
Rump color conditions, genetic analysis 496
Rump color inheritance due to interaction of factors 497
Rump colors
Blue 494, 495
Black 495, 497
Dark blue 495, 497
Light blue 495
Silvery blue 495
Uniform blue 495
Recessive 496
White 494, 495, 497
Dominant to blue 496
Segregation 475
Selection experiment 468, 484–486
Self-black Tumbler 500
Sooty 469, 471, 472, 483, 485
Species 467 (See Columba)
Differences based on single factors 505
Spot 466
Tumblers 482
Typical blue or silver tail 498

Pigeons (*continued*)
Variation in color of outer vanes of outer tail feathers 487–494
White 467, 486, 487
White spotting 469
WHITMAN'S results 469
PLATE, L. 583, *589*
PLOUGH, H. H. *334*, 338, 373, *380*
Pollen grains in parents and F_1 hybrids in wheat 523–526
Pollen grains segregation in F_1 rice hybrid 545
Pollen of F_1 plants of species hybrids 545
Polymorphic species of domesticated plants and animals 423, 425
Polyphyletic origin domesticated pigeon 505
Polyploid species chromosome behavior 531–533
Polyploid species crosses between in wheat 530
Polyploid species genetic behavior 527–530
Polyploid species results of crossing 529
Polyploid species of wheat sterility 533
Polyploidy correlation with multiple factors 540
Polyploidy in plant species 526, 527
POPE, M. N. 239
Presence and absence theory 424
Primula kewensis 526
Primula sinensis 597
Primula tetraploid 528, 530
Primula flower color 333
Protein content of maize 238
Protozoa 145
Pure line hypothesis still valid 181
Pure line problem 175
Pure lines, results of selection in *Pestalozzia Guepini* (which see) 142–201
Pygaera curtula L.×*P. anachoreta* F. F_1 532

QUINCKE, HENRICUS 574, *582*

Random mating 508–511
Ratios modified in wheat 540
Rats, white
Alcoholism and growth 427–445
Alcohol as selective agent 134
Alcohol effect on number and size of litters 117
Alcohol treatment reduces number of young produced 139
Drunken by inhalation 123, 140
Fertility 125
Criteria of influence of alcohol treatment 124
Greater in tests than in controls in later generations 134
Germ cells inferior eliminated by alcohol 134
Germplasm modification due to alcohol 118, 139, 140
Infertility eliminated by alcohol 118
Litter number decreased by alcohol 140
Litter size 126
Increased in treated males 136
Of treated rats reduced 129, 130, 132, 135, 140, 141
Litters number compared in guinea pig and rat treated with alcohol 137
Litters of treated rats increased number in later generations 134
Modification due to alcohol 141
Oestrous cycle lengthened by alcohol treatment 137
Rats, offspring of alcohol treated genetically superior to controls 441
Rats, resistance to effect of alcohol 136
Selective elimination due to alcohol 139–141
Treatment of rats 122
WISTAR defectives 124, 127
RAY, JOHN *507*
Recessive factors effective in presence of dominant 545
Recessive factors in excess may be dominant over normally dominant factors 527
Recessive genes two dominate the normal 527
Recombination, origin of characters by 422
Reduction divisions in wheat and rye 548
Reduplicated chromosomes 527
RHODE ISLAND AGRICULTURAL EXPERIMENT STATION 469
RHODE ISLAND RED breed of fowl 446–465, 447, 448, 452, 463
RICHARDS, M. H. *380*
RIDDLE, OSCAR *507*
RIDGWAY 472
ROBERTS, ELMER 333, *334*
ROBBINS, RAINARD B. 508–512
ROOT, F. M. 143, 144, 175, 176, 181–*183*
Rosa, chromosome number 526, 531–534
Rosa hybrids 532
Rosa Sabini (*R. pimpinellifolia*×*R. sylvestris*) 532
ROSENBERG, O. 526, 531, 535, *551*, *552*
ROSSI, GILBERTO 584
RÜSSOW 569
Rust resistance in wheat 514
Rye chromosomes 519, 547

SACHS 569
SAKAMURA, T. 514, 515, 517, 543, *552*
SALVADORI, T. 467, 495, 505, 507
SAX, KARL 513–558, 514, 536, *552*
SCHACHTZABEL E. 473, *507*
SCHLOSS, OSCAR M. 573, *582*
SCHOFIELD, RICHARD 583, *589*
Secale cereale 547
Seeds 269
Number of ovules per pod 269, 271
Number per pod 269, 271
Pods per plant 271
Seeds treatment to secure complete germination 592
Weights 269, 271
Segregation in pollen grains in F_1 rice hybrid 545
Selecting device for isolating spores 168
Selection
According to progeny character 176
As factor in evolution 175

Selections (*continued*)
By elimination of hereditary factors in Drosophila 113
Choice of seeds from dominant plants 512
Effect of evidence wanting 181
Effect of in Drosophila 114
Nature of 113
Effectiveness in bar eye Drosophila 112
Effects conflicting evidence 143, 144
Experiments in maize 237
Experiments in pigeons 468, 484–486
Factors influencing results of 174
For eye facet number in *Drosophila melanogaster* 1–115
For length of spores **158**–162
For style length and fertility of *Oenothera brevistylis* **590**–596
Germinal changes during course of 2, 78
In an allogamous population **508**–512
In Arcella 176, 178, 179, 181
In Centropyxis 176, 181
In Didynium 176, 181
In Difflugia 178, 179, 181, 182
In Paramecium 182
In Stylonychia 144, 175, 176
Ineffective 143
Mass results 176
Method 180
Of spores according to visible characters **167**–174
Positive results by conserving mutations 4
Possibilities of 237
Result negative 174
Results 177–180
In Pestalozzia 175
With Difflugia 144
Within pure lines of *Pestalozzia Guepini* (which see) **142**–201
Selections eliminating germinal factors, effective 112
Selective effect of alcohol on rats 440, 441
Selective elimination due to alcohol 139–141
Self-fertilization producing disease resistant races in maize 252
Self-fertilized lines in maize improvement **252**–257
Sex character, secondary 583
Sex-linked characters in Drosophila 583
Sex-linked characters in *Drosophila obscura* **335**–384
Sex-linked characters in man 583
None behaves as sex-linked characters in all families 588
Three principles 588
Sex-linked groups of Drosophila species compared 373
Sex-linked heredity 583
Sex-linked mutant for *Drosophila repleta* 320
SEYSTER, E. W. 3, 8, 16, *115*
SHAMEL, A. D. 543, *552*
SHEPPARD'S tables 156
SHULL, G. H. 253, 530, *552*
Ski not a monogenic character 399
Ski wings in *Drosophila melanogaster* **385**–426
SMALLEY, F. W. 468, 475–477, *506*
SNYDER, LAURENCE H. 427
Somatic development and parental sets of chromosomes in F_2 hybrids 558
Sorghum vulgare glutinosum 569
Species crosses in wheat 553
Species differences in pigeons based on single factor 505
Species groups in wheat antiquity of 541
Species hybrids in animals chromosomes behavior in F_1 532
Species hybrids in wheat **513**–552
Behavior of F_1 chromosomes in reduction division 515, 517, 518
Chromatic bodies in microspores 515
Chromosome number 525
Chromosomes in sterile and partially sterile **520**–521
Correlation between chromosome number and fertility in F_2 515
Sterile **535**
Variation in chromosome number 515
Species polymorphic wild and domesticated 425
Species polyploid genetic behavior **527**–530
SPILLMAN, W. J. 514, *552*
STAPLES-BROWNE, R. 468, 471, 472, 475–477, 480, 486, 496, 497, 507
Sterile F_2 due to vegetative development and gametic sterility 558
Sterility and chromosome relationships 553
Sterility and quantitative relations between univalents and trivalents 536, 537
Sterility and reduplication of single chromosome 534
Sterility in wheat hybrids **513**–558
Cause 536, 537, 540, 549
STEVENS, F. L. 182, *183*
STOCKARD, C. R. 122–124, 126, 131, 134–136 139, *141*, 428, 442, *445*
STOCKING, RUTH J. 144, 175, *183*
STURTEVANT, A. H. 319–321, 333, *334*, 337, 338, 371, 373, 376, 378, *380*, 386, 391, 397, 423, *426*, 535, *552*
Stylonychia 144, 175, 176
Response to selection 175
SURFACE, F. 274, 276, 297, *318*
Swertia perennis 569

TÄCKHOLM, G. 526, 531, *552*
TAHARA, M. 527, *552*
TEGETMEIER, W. B. 467, *507*
Temperature effect on inheritance of flower color 333
Tetany 581
Tetraploid forms 526, 527
Of tomato and nightshade 543
Originated by chromosome duplication 540
Reduction 540
Tetraploidy in wheat species origin **540**–543
Tetraploidy value **543**, 545
THATCHER, L. E. 508
THOMPSON, D. H. 1, 70, *115*
TICE, S. A. 5, *115*
TREUB 569

Triploid hybrids 533
Triploid individual 534
Triticum (see wheat)
Triticum aegilopoides 542
Triticum aegilops 516
Triticum aegilops cylindrica 541
Triticum aegilops ovata 525, 541
Triticum aestivum subsp. *durum* 516
Triticum aestivum subsp. *vulgare* var. *compactum* 516
Triticum aestivum subsp. *vulgare* var. *Spelta* 516
Triticum aestivum subsp. *vulgare* var. *turgidum* 516
Triticum dicoccoides 541, 542
Triticum sativum, *T. aestivum* subsp. *vulgare* var. *erythrospermum* (Preston) Körn *albidum* Körn (Amby) (Bluestem) 516
Sterility in pentaploid forms 534
Einkorn group 513, 516, 517, 525, 526, 534, 541, 542
Triticum monoccocum 513, 514, 516, 517, 542, 544, 546
Var. *Hornemanni* 516
Emmer group 513, 516, 518, 519, 525, 526, 534, 537, 539, 540–542, 544, 547
Triticum dicoccum 513, 514, 518, 542
Triticum durum 513–516, 518, 542, 553
Triticum polonicum 513, 514, 516, 518, 542
Triticum pyrmidale 542
Triticum turgidum 513, 514, 516, 518, 542, 546
Vulgare group 513, 516, 518, 519, 525, 526, 534, 537–542, 544, 547
Triticum compactum 513, 514, 516, 542
Hybrid origin of 541, 543
Triticum Spelta 513, 514, 516, 518, 542
Triticum sphaerococcum 542
Triticum vulgare 513, 514, 516, 519, 525, 542, 553
Sterility in pentoploid forms 534
TSCHERMAK, E. VON 514, *552*
TUPPER, W. W. 545, *552*
TURPIN, C. 275, 277, *317*
Turtur lugens 502
Turtur turtur 502

UNIVERSITY OF ILLINOIS 1
UNIVERSITY OF MICHIGAN 142, 590
UNIVERSITY OF MISSOURI 559
UNIVERSITY OF WISCONSIN 466
Urticaria 575, 581
UTAH AGRICULTURAL EXPERIMENT STATION 277

VALENTI, ANNA LOUISA 588, *589*
VANDER VEER, ALBERT JR. 576, *581*
VAVILOV, N. I. 514, *552*
VICARI, E. M. 117, 122, *141*, 427, 435, *444*
DE VRIES 528

WALDECK, MRS. 506
WALKER, ISAAC CHANDLER 574, *582*
WARNER, D. E. 275, *317*, 446, 447, 452, 464, *465*
WEATHERWAX, PAUL 203, *226*, 568–572
WEINSTEIN, ALEXANDER 419, *426*
WENE, ELMER H. 277, *317*
Wheat
Assortment of chromosomes 540
Bud mutations associated with tetraploidy 543
Cause of sterility in wheat hybrids 536, 540, 549
Cell division in species 544, 546, 548
Chromatin granules outside of nuclei in pollen grains 521
Chromosome behavior and origin of species 514
Chromosome duplication 550
Chromosome linkage 539
Chromosome number and adaptability in species 542
Chromosome number and cell size correlated 519, 523, 540
Chromosome number and size of pollen grains 523–526, 549
Chromosome number and sterility 534
Chromosome number in species 514, 525, 540, 541–543, 547, 553, 557, 558
Chromosome reduplication 540
Chromosome complete set necessary for gametic and somatic development 537
Chromosomes in sterile and partially sterile hybrids 520–521
Chromosomes of species 514, 517–521
Correlation between chromosome number and fertility in F_2 515
Correlation between sterility and vegetative development 535
Correlation between grain weight and sterility 557
Correlations in hybrid sterility in wheat 514
Cytological methods 517
Endosperm 553, 554, 556, 558
Endosperm development and sterility 553–558
Endosperm formation not due to triploid condition 558
Grain 555
Variability and chromosome numbers 558
Heterozygosis and pollen variability 525–526
Hexaploidy origin 540, 541
Hybrids 513, 514, 516, 520–525, 530, 531, 534, 535, 538, 539, 546, 547, 549, 554, 557
Intermediate chromosome numbers absent 549
Linked groups 538
Modified ratios 540
Pollen grains in parents and F_1 hybrids 523–526
Polyploid species sterility 533
Polyploidy in plant species 526–527
Reduction divisions 548
Seeds weight and sterility 553, 554
Segregates 549
Somatic development and parental chromosomes in hybrids 558
Species groups, antiquity 541

Wheat (*continued*)
Species, group differences 541
Sterility 536, 538, 549, 555
Sterility and chromosome relationships 553
Sterility and vegetative development 557
Sterility groups 525, 541
Sterility in hybrids **513**–558
Sterility of species hybrids 549
Tetraploidy origin **540**, 541
Variability 542
Variation in chromosome number 515
White Leghorn breed of domestic fowl, 446–448, 451, 452, 454, 463, 465
WHITMAN, C. O. 468, 472, 476, 477, 484, 486, 496, 505–*507*
WHITMAN'S results interpreted on genetic basis **469**
WHITMAN'S selection experiments 504
WIEMAN, H. L. 587, *589*
WILCOX, A. N. 247
WILLIAMS, C. G. 508
WILLOUGHBY, F. 466, *507*
WILSON, E. B. *380*
WINKLER, H. 543, *552*
WINTON 239
WISCONSIN AGRICULTURAL EXPERIMENT STATION 469, 474
WISTAR INSTITUTE 118, 124, 127
WOLLENWEBER, H. W. 150, *183*
WRIGHT, SEWALL 258, 261–265, 267–269, 271, *273*

YAMAGUCHI, YASUKÉ 569, *572*
YULE 263

ZADE *552*
Zea mays, see maize
ZELENY, CHARLES 1–115, 2, 4, 6–8, 13, 19, 31, 84, *115*
Zenaidura macroura carolinensis 502
Zygotes eliminated by alcohol 140